

# Anagrain

Metoclopramide hydrochloride / Paracetamol

## Consumer Medicine Information (CMI)

### What is in this leaflet

This leaflet answers some common questions about Anagrain. It does not contain all the available information. It does not take the place of talking to your pharmacist or doctor.

All medicines have risks and benefits.

**If you have any concerns about taking this medicine, talk to your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

### What Anagrain is used for

Anagrain is used for the relief of headache, nausea and vomiting associated with migraine.

Anagrain contains the active ingredients metoclopramide (as metoclopramide hydrochloride anhydrous) and paracetamol.

Metoclopramide helps control nausea and vomiting caused by migraine and other illnesses. It works by blocking the action of a chemical in the brain which causes nausea and vomiting. It also acts in the stomach and upper intestine to increase muscle contractions.

Paracetamol is an analgesic. It provides effective temporary relief from pain.

There is no evidence that Anagrain is addictive.

### Before you take it

#### *When you must not take it*

**Do not take Anagrain if you have ever had an allergic reaction to:**

- metoclopramide
- paracetamol
- any of the ingredients listed at the end of this leaflet.

Symptoms of an allergic reaction may include asthma, wheezing, shortness of breath, swelling of the face, lips or tongue which may cause difficulty in swallowing or breathing, skin rash, itching or hives.

**Do not take it if you have any of the following conditions:**

- bleeding from the stomach and/or digestive tract
- intestinal blockage
- recent surgery on the stomach and/or digestive tract
- phaeochromocytoma (a rare tumour of the adrenal gland).

**Do not take Anagrain if:**

- **you are pregnant, or intend to become pregnant**

It may affect your developing baby if you take it during pregnancy.

- **you are breast-feeding or intend to breast-feed**

Anagrain passes into breast milk and therefore may harm the baby.

**Do not take Anagrain after the expiry date (EXP) printed on the pack.**

**Do not take it if the packaging is torn or shows signs of tampering or if the tablets do not look quite right.**

**Do not give this medicine to children under 12 years of age.**

The safety of this medicine in children under 12 years of age has not been established.

**Talk to your doctor or pharmacist if you are not sure whether you should start taking Anagrain.**

#### *Before you start to take it*

**Tell your doctor or pharmacist if:**

1. **you have allergies to any other medicines, foods, preservatives or dyes.**
2. **you have or have had any of the following medical conditions:**
  - epilepsy - metoclopramide may increase the risk of you having a fit
  - liver or kidney disease
  - Parkinson's disease - metoclopramide may make this condition worse.

**If you have not told your doctor or pharmacist about any of the above, tell them before you take Anagrain.**

#### *Taking other medicines*

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.**

Some medicines may be affected by Anagrain or may affect how well it works. You may need to take different amounts of your medicines or you may need to take different medicines. Your doctor or pharmacist has a more complete list of medicines to avoid while taking Anagrain.

---

## How to take it

---

### *How much to take*

Anagrain should be taken at the first sign of a migraine attack.

#### **Adults:**

The initial dosage for adults is 1-2 tablets and then 1-2 tablets every four hours, as needed. **Do not take more than 6 tablets in 24 hours.**

#### **Adolescents (12-17 years):**

The initial dosage for adolescents 12 to 17 years old is 1 tablet and then 1 tablet every four hours, as needed. **Do not exceed 3 tablets in 24 hours.**

**Do not take Anagrain for longer than 48 hours at a time unless advised to by a doctor.**

### *How to take it*

**Swallow the tablet with a glass of water.**

### *If you forget to take it*

**If symptoms persist take your next dose when you are meant to.**

**Do not take a double dose to make up for the dose that you missed.**

This may increase the chance of you getting an unwanted side effect.

**If you are not sure what to do, ask your doctor or pharmacist.**

**If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.**

### *If you take too much (overdose)*

**Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Anagrain. Do this even if there are no signs of discomfort or poisoning.**

You may need urgent medical attention.

If you take too much Anagrain you may experience the following symptoms:

- vomiting
- stomach pain
- sweating
- low blood pressure
- yellowing of the skin
- drowsiness
- confusion
- twitching or uncontrolled spasms.

---

## While you are taking it

---

### *Things you must do*

**Tell your doctor or pharmacist if nausea, vomiting or headache persists.**

**Tell all doctors, dentists and pharmacists who are treating you that you are taking Anagrain.**

**If you are about to be started on any new medicine, tell your doctor or pharmacist that you are taking Anagrain.**

**Tell your doctor or pharmacist immediately if you become pregnant while taking Anagrain.**

### *Things you must not do*

**Do not take Anagrain for longer than 48 hours at a time unless advised to by a doctor.**

**Do not take Anagrain to treat any other complaints unless your doctor or pharmacist tells you to.**

### *Things to be careful of*

**Do not take Anagrain with other products containing paracetamol, unless advised to do so by a doctor or pharmacist.**

Not more than 4 g of paracetamol should be taken in any 24 hour period.

**Be careful driving or operating machinery until you know how Anagrain affects you.**

---

## Side effects

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Anagrain.**

Anagrain helps most people with migraines but it may have some unwanted side effects in some people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

**Tell your doctor or pharmacist if you notice any of the following common side effects and they worry you:**

- drowsiness, tiredness, dizziness, restlessness, fatigue
- bowel upsets
- upset stomach
- allergic reactions.

Other side effects not listed above may also occur in some people.

Ask your doctor or pharmacist to answer any questions you may have.

---

## After taking it

---

### *Storage*

**Keep your tablets in the blister pack until it is time to take them.**

If you take the tablets out of the blister they may not keep as well.

**Keep your tablets in a cool dry place where the temperature stays below 30°C.**

**Do not store Anagrain or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.**

Heat and dampness can destroy some medicines.

**Keep it where young children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### *Disposal*

**Dispose of the tablets where children cannot reach them.**

**If you stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.**

---

## Product description

---

### *What it looks like*

Anagrain tablets are round, white and scored on one side. They are available in packs of 8 tablets.

### *Ingredients*

#### **Active ingredients:**

Each tablet contains 5 mg of metoclopramide hydrochloride anhydrous and 500 mg of paracetamol.

#### **Inactive ingredients:**

- magnesium stearate
- sodium starch glycollate.

Anagrain tablets do not contain lactose, gluten, tartrazine or any other azo dyes.

---

## Sponsor

---

Aspen Pharmacare Australia Pty Limited  
34-36 Chandos St  
St Leonards NSW 2065

Australian Registration Number:  
AUST R 13547.

This leaflet was prepared in February 2010.